

Durham Research Online

Deposited in DRO:

02 February 2012

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Sawdon, M. and Finn, G. and Griksaitis, M. and Kirkman, E. (2010) 'Simulating haemorrhage in medical students using a lower body negative pressure chamber.', Association for Simulated Practice in Healthcare (ASPiH) 2010 Inaugural Annual Conference. Newcastle-upon-Tyne, England, 17-18 November 2010.

Further information on publisher's website:

<http://www.aspih.org.uk/>

Publisher's copyright statement:

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

**Title: Simulating Haemorrhage in Medical Students using a Lower
Body Negative Pressure Chamber**

Authors:

Dr Marina Sawdon BSc (Hons) PhD

Lecturer in physiology, School of Medicine & Health

Durham University, Queen's Campus

Ms Gabrielle Finn BSc (Hons)

Lecturer in anatomy, School of Medicine & Health

Durham University, Queen's Campus

Dr Emrys Kirkman BSc (Hons) PhD

Honorary Senior Lecturer, School of Medicine & Health

Durham University, Queen's Campus

Dr Michael Griksaitis MBBS, FRCP

Clinician, Neonatal Intensive Care Unit

James Cook University Hospital,

Correspondence:

Dr Marina Sawdon

Phase 1 Medicine

University of Durham, Queen's Campus

University Boulevard

Stockton-on-Tees

Cleveland

TS17 6BH

Email marina.sawdon@durham.ac.uk

Tel +44 191 3340340

Fax +44 191 3340321

KEYWORDS:

Doppler ultrasound

Haemorrhage

Simulation

Stroke volume

Medical education

Abstract

Introduction This article is a descriptive report of a novel way of teaching the cardiovascular response to progressive haemorrhage in a first year medical undergraduate setting using simulation. Simulation may provide the means to allow students to see in practice the theoretical knowledge they have gained from lectures, thus giving clinical relevance to that knowledge.

Method The early phase (compensatory phase 1) of a 'simple' progressive haemorrhage is simulated in 6 volunteer medical students by applying sub-atmospheric pressure to the lower body using a lower body negative pressure (LBNP) chamber (see figure below). This is a technique that has been widely used in research to simulate

orthostatic stress and haemorrhage. The sub-atmospheric pressure causes 'pooling' of blood in the vessels (particularly veins) of the legs and pelvis. This 'pooling' reduces the amount of blood returning to the heart (venous return) and thus mimics the effects of losing blood from the cardiovascular system. Parts of this method have been described elsewhere (1). To demonstrate the physiological responses to LBNP a range of cardiovascular parameters are monitored throughout the demonstration to allow students to observe the integrated (whole body) response to 'progressive haemorrhage'. Cardiac stroke volume is monitored using a portable ultrasound device (SonoSite MicroMaxx™), heart rate is measured accurately from an ECG trace and blood pressure using an automated sphygmomanometer. Tissue (skeletal muscle) oxygenation which, in this context reflects changes in blood flow in the deltoid, forearm and thenar eminence muscles is assessed using Near Infrared Spectroscopy (NIRS).

Results Progressive haemorrhage will reduce venous return and preload, thus reducing stroke volume by Starling's law of the heart (2) and ultimately reduce arterial pulse pressure. The body's initial response (mediated by the baroreceptor reflex) to the reduced arterial pulse pressure is an increase in heart rate to maintain mean arterial blood pressure (3) in the face of a falling cardiac output. The students are required to calculate the changes in overall total peripheral resistance and are able to see the consequences of the resulting reduction in muscle blood flow as a fall in muscle oxygenation.

Conclusion Durham University Medical Programme uses simulation in a novel way to reinforce didactic teaching of the cardiovascular response to haemorrhage utilising a

LBNP chamber. The simulated haemorrhage may provide the means to allow students to see in a 'clinical' context the theoretical knowledge they have gained from lectures, which may improve knowledge retention.

Word count 397

References

1. Sawdon M & Kirkman E. Simulating Haemorrhage in Medical Students. *International Journal of Clinical Skills*. 2008; **2**(2): 74-78.
2. Patterson SW, Starling EH. On the mechanical factors which determine the output of the ventricles. *Journal of Physiology*. 1914; **48**(5): 357-379.
3. Sawdon M, Ohnishi M, Watkins P & Kirkman E. The effects of primary thoracic blast injury and morphine on the response to haemorrhage in the anaesthetised rat. *Experimental Physiology*. 2002; **87**(6): 683-689.

